


Investigation of rate constant of Fenton-like reactions in the presence of Glutamic and Glutaric acid at pH 5 determined by UV/VIS spectroscopy

Nazanin Taherkhani, Daniele Scheres Firak, Thomas Schaefer, and Hartmut Herrmann

Leibniz Institute for Tropospheric Research, Atmospheric Chemistry Department (ACD), Leipzig, Germany

Contact: taherkhani@tropos.de



Introduction

- Fenton reactions are crucial in producing $\cdot\text{OH}$ in the atmospheric aqueous particle phase.
- Reported second-order rate constants for **Fenton** ($\text{Fe}^{2+} + \text{H}_2\text{O}_2$) at $68 \text{ L mol}^{-1} \text{ s}^{-1}$. This rate can be modified with the Fe speciation (**Fenton-like**).^[1]
- Glutamic acid**, as a multifunctional amino acid which is abundant in nature, undergoes different protonation states depending on the pH of the solution^[2]
- Iron can form a complex with **amino acids**, such as glutamic acid, which may significantly modify the reaction kinetics.





Fig.1: Proportional distribution of amino acids in the coarse-mode particles as combined amino acids. ^[1]



Experimental

Iron conversion

- APHA 3500 Fe test (APHA, 1997).^[3]
- Fe(II) and ortho-phenanthroline complex.
- DMSO used as a scavenger of $\text{HO}\cdot$

Rate constants determination

- First and second-order rate constant.
- Employing UV-Vis spectroscopy.

Redox Potential

- Explore the redox behavior of the $\text{Fe}^{2+} / \text{Fe}^{3+}$ in the presence of glutamic acid
- DMSO used as a scavenger of $\text{HO}\cdot$




Fig. 2: Formation of complex between Fe(II) and 1,10-Phenanthroline




Fig. 3: Absorption spectra of the complex at $4.5 \times 10^{-5} \text{ mol L}^{-1} \text{ Fe}$ and $4 \times 10^{-4} \text{ mol L}^{-1} \text{ H}_2\text{O}_2$ throughout the Fenton reaction

Results

Kinetics of Fenton-like reaction




Fig. 4: Speciation of Fe(II) in the presence of Glutamic acid at different pH. Conc. Fe(II) $4 \times 10^{-3} \text{ mol L}^{-1}$ and Glutamic acid $4 \times 10^{-4} \text{ mol L}^{-1}$. (Visual MINTEQ 3.1)

- Low pH favors the formation of certain Fe–glutamate complexes.
- As pH increases, the speciation of these complexes shifts, influenced by changes in ligand protonation and metal–ligand binding affinities (availability of free Fe^{2+} ions decrease) leading to the formation of insoluble Fe(OH)_2 or the oxidation to Fe^{3+} ions, which form Fe(OH)_3 precipitates^[4]

First-order and second-order rate constant




Fig.5 : Plot of $\ln$ of Fe(II) concentration as a function of time at different concentrations of hydrogen peroxide for glutamic acid

- The first-order rate constant was calculated by analyzing the reaction progress at different H_2O_2 concentrations. The reaction rate was proportional to the concentration of H_2O_2 which ranged from 1.5×10^{-4} to $6 \times 10^{-4} \text{ mol L}^{-1}$.
- The presence of the ligand significantly increased the slope of the plots, leading to accelerated the reaction rates.
- The slope with glutamic acid was higher than without any ligand, indicating a noticeable effect of complexation of glutamic acid with iron on the kinetics of the reaction.




Fig.6 : Plot of the first-order rate constants against the different hydrogen peroxide for glutamic acid, glutaric acid and without ligand.

Slope corresponds to the **second-order rate constant** for the **Fenton-like** reaction

Table 1. Second-order rate constants obtained with different ligands at pH 5

Reactions at pH 5	$k_2 / \text{L mol}^{-1} \text{ s}^{-1}$
without ligand	68 ± 2
with glutamic acid	102 ± 5
with glutaric acid	82 ± 2
with glutamic acid (observed)	130

Conclusions

- The Fenton-like reaction at pH 5 was initially measured with a second-order rate constant of $68 \pm 2 \text{ L mol}^{-1} \text{ s}^{-1}$ in the absence of any ligand, serving as the reference.
- By introducing glutamic acid, the rate constant increased to $102 \pm 5 \text{ L mol}^{-1} \text{ s}^{-1}$, indicating a significant enhancement in reaction kinetics due to ligand coordination.
- Glutaric acid increased the second-order rate constant ($82 \pm 2 \text{ L mol}^{-1} \text{ s}^{-1}$) compared to the reaction without a ligand, which was less significant compared to glutamic acid.
- The Fe-glutamic acid complex, with a redox potential of 0.514V, exhibits favorable thermodynamic properties, highlighting the stability and reactivity of iron-glutamic acid complexes in redox processes.

Improving Complex Formation with UV-vis Spectrum Analysis




Fig. 7: UV-Vis Spectrum for Forming complexes with two different concentrations of glutaric acid at pH 5




Fig. 8: Formation of the complex using structural formulas

- The UV-vis spectrum shows iron (Fe^{2+}) forms a stable complex with glutaric acid. This shift to lower wavelength indicates coordination between Fe^{2+} and glutaric acid, similar to glutamic acid complexes.
- While the stability constant for the Fe^{2+} -glutamic acid complex is known, the value for the Fe^{2+} -glutaric acid complex is unspecified. Based on the similarity between the two complexes, a comparative spectral investigation was conducted to explore their complexation behavior.

Redox Potential




Fig. 9: Redox reactivity of organically complexed iron(II) species ^[6]

- Glutamic acid influences the redox behavior of Fe ions by altering their oxidation potentials.
- Stable complexes are formed with Fe^{2+} and Fe^{3+} , affecting their electrochemical properties, and the redox potential of the $\text{Fe}^{2+}/\text{Fe}^{3+}$ gained from Nernst equation:

$$E = E^\circ - \frac{RT}{nF} \ln Q \quad [5]$$

- The redox potential of 0.514 V demonstrates the important role of ligands like glutamic acid in modifying iron's electrochemical properties.

Atmospheric aqueous phase implications

- In the atmospheric aqueous phase, organic acids such as glutamic acid can complex with metal ions like Fe^{2+} , modifying their reactivity and influencing atmospheric chemistry. This metal-ligand complexation is critical as it regulates the formation of reactive oxygen species (ROS), particularly hydroxyl radicals ($\text{HO}\cdot$), which are key to the oxidation of pollutants and aerosols.^[7]
- These complexes are particularly important in aerosols and cloud droplets, where they facilitate redox reactions that drive atmospheric oxidation cycles under acidic conditions (pH 1-5). ^[8]

References

- [1] Koppenol, Dalton Trans., **2022**, 51, 17496-17502. [2] Tanaka, Y. & Nakamura, K., **2023**, Env. Sci. Technol. 57 (10), 1234-1245. [3] APHA (**1997**) Standard Methods for the examination of water and wastewater Visual MINTEQ 3.1 .[4] Visual MINTEQ (3.1) [5] Feiner A.S. & McEvoy A.J.,**1994**, J. Chem. Educ., 71(6), 493. [6] Strathmann T. J., **2011**, Redox reactivity of organically complexed iron(II) species with aquatic contaminants. [7] Tilgner et al., Atmos. Chem. Phys., **2021**, 21,p.13483-13536. [8] Song et al., RSC Adv., **2024**, 14, 33489-33511.