

ORAL PRESENTATION

Advanced modelling of the multiphase DMS chemistry with the CAPRAM DMS module 1.0

E. H. Hoffmann¹, A. Tilgner¹, R. Schrödner^{1,2}, P. Bräuer^{1,3}, R. Wolke¹, H. Herrmann¹. (1) Leibniz Institute for Tropospheric Research (TROPOS), Permoserstraße 15, 04318 Leipzig, Germany. (2) Centre for Environmental and Climate Research, Lund University, SE-22362 Lund, Sweden. (3) Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York YO10 5DD, United Kingdom..

Oceans are the general emitter of dimethyl sulphide (DMS), the major natural sulphur source [1], and cover approximately 70 % of earth's surface. The main DMS oxidation products are SO₂, H₂SO₄ and methyl sulfonic acid (MSA). Hence, DMS is very important for formation of non-sea salt sulphate (nss SO₄²⁻) aerosols and secondary particulate matter and thus global climate. Despite many previous model studies, there are still important knowledge gaps, especially in aqueous phase DMS chemistry, of its atmospheric fate [2].

Therefore, a comprehensive multiphase DMS chemistry mechanism, the CAPRAM DMS module 1.0 (DM1.0), has been developed. The DM1.0 includes 103 gas phase reactions, 5 phase transfers and 54 aqueous phase reactions. It was coupled with the multiphase chemistry mechanism MCMv3.2/CAPRAM4.0a ([3]; [4]) and the extended CAPRAM halogen module 2.1 (HM2.1, [5]) for investigation of multiphase DMS oxidation in the marine boundary layer. Then, a pristine ocean scenario was simulated using the air parcel model SPACCIM [6] including 8 non-permanent cloud passages - 4 at noon and 4 at midnight. This allows the investigation of the influence of deliquesced particles and clouds on multiphase DMS chemistry during both daytime and nighttime conditions as well as under cloud formation and evaporation. To test the influence of various subsystems on multiphase DMS chemistry different sensitivity runs were performed. Investigations of multiphase chemistry of DMS and its important oxidation products were done using concentration-time profiles and detailed time-resolved reaction flux analyses.

The model studies revealed the importance of aqueous phase chemistry for DMS and its oxidation products. Overall about 7.0% of DMS is effectively oxidised by O₃ in the aqueous phase of clouds. The simulations revealed the importance of halogen and aqueous phase chemistry for DMS and its oxidation products. Overall halogen compounds contribute with 71% to DMS oxidation with gaseous Cl (18.4%) and BrO (46.1%) as main oxidants. The conversion efficiency of DMS to SO₂ in the gas phase was simulated between 0.24, 0.31 and 0.6 for the full pristine ocean scenario run, a simulation without considered halogen chemistry and a simulation without treated aqueous phase DMS chemistry, respectively. Furthermore, the studies indicate that the conversion efficiency of DMS to MSA is strongly related to DMS oxidation by BrO and treating of aqueous-phase DMS chemistry. The MSA yield for different sensitivity runs was simulated between 0.01 and 0.47. The lowest yield is reached treating only gas phase chemistry of DMS. Moreover, the simulation with the whole mechanism indicate that multiphase DMS oxidation produce as much MSA as sulphate leading to strong implications for nss-SO₄²⁻ aerosol formation, activation to cloud condensation nuclei and cloud albedo.

References

- [1] Andreae, M. O., Mar. Chem., 1990, 30, 1-29.
- [2] Barnes, I., et al., Chem. Rev., 106, 940-975, 2006.
- [3] Rickard, A., et al., The Master Chemical Mechanism Version MCM v3.2, available at: <http://mcm.leeds.ac.uk/MCMv3.2/> (last access: 05 Mai 2015), 2015.
- [4] Bräuer, P., et al., Atmos. Chem. Phys. Discuss., in preparation, 2017.
- [5] Bräuer, P., et al., J. Atmos. Chem., 70, 19-52, 2013.
- [6] Wolke, R., et al., Atmos. Environ., 39, 4375-4388, 2005.